

STUDENT \_\_\_\_\_

CLASS \_\_\_\_\_

## LABOR, CHILDBIRTH, AND POSTPARTUM TERMINOLOGY

Match the correct number with the definition shown below.

ACTIVE LABOR  
 AFTERBIRTH  
 ANALGESIC  
 ANESTHETIC  
 ANESTHESIOLOGIST  
 APGAR SCORE  
 BRAXTON-HICKS CONTRACTIONS  
 BREECH  
 CESAREAN  
 CAUDAL BLOCK  
 CIRCUMCISION  
 COLOSTRUM  
 CONTRACTIONS  
 CROWNING  
 DELIVERY  
 DILATION  
 EFFACEMENT  
 ENEMA  
 EPIDERMAL BLOCK  
 EPISIOTOMY  
 FONTANEL  
 GYNECOLOGIST  
 INACTIVE LABOR

INDUCED ABORTION  
 INVOLUTION  
 LABOR  
 LIGHTENING  
 LOCHIA  
 MISCARRIAGE  
 NATURAL ABORTION  
 NATURAL CHILDBIRTH  
 OBSTETRICIAN  
 PARICERVICAL BLOCK  
 PEDIATRICIAN  
 PERINEUM  
 PLACENTA  
 POSTNATAL  
 POSTPARTUM  
 PREMATURE  
 PUDENDAL BLOCK  
 SADDLE BLOCK  
 SHOW  
 SPINAL BLOCK  
 STATION  
 STILL BIRTH  
 TRANSITION

1. \_\_\_\_\_ when the uterus contracts at irregular intervals throughout pregnancy. They exercise and prepare the uterus for labor and help circulate maternal blood.
2. \_\_\_\_\_ when a full-term baby is dead at birth.
3. \_\_\_\_\_ low spinal, numbs from lower belly down, must stay flat 18-24 hours, and often be catheterized.
4. \_\_\_\_\_ the soft spot on top of the baby's head where the skull bones have not grown together.
5. \_\_\_\_\_ when a baby is delivered through a cut in the abdominal wall.

6. \_\_\_\_\_ labor from 8-10 centimeters dilation, often considered the hardest part of labor. It ends the first stage of labor.
7. \_\_\_\_\_ anesthetic given in cervix and lower lateral border of uterus to numb that area.
8. \_\_\_\_\_ rich, watery fluid that comes from mother's breast before milk comes.
9. \_\_\_\_\_ a procedure that terminates the fetus before full term.
10. \_\_\_\_\_ childbirth without need of pain killers; preparation to relieve fear, tension, and pain before birth.
11. \_\_\_\_\_ refers to the level of the infant's head as it descends into the birth canal.
12. \_\_\_\_\_ an injection between vertebrae in the lower back to numb the navel to thigh area.
13. \_\_\_\_\_ the placenta, umbilical cord, amniotic sac, and lochia delivered as the third stage of labor.
14. \_\_\_\_\_ anesthetic given in spinal fluid around the spinal cord.
15. \_\_\_\_\_ time from birth to 6 weeks for mother.
16. \_\_\_\_\_ birth of the baby from birth canal to outside world.
17. \_\_\_\_\_ a physician who is trained in female concerns.
18. \_\_\_\_\_ the purpose is to open the cervix to about 3 cm. It is the longest and least uncomfortable labor and often occurs at home.
19. \_\_\_\_\_ time from birth to 6 weeks for baby.
20. \_\_\_\_\_ the plug covering the cervix. It is released at the beginning of labor and is pinkish red in color.
21. \_\_\_\_\_ a drug that relieves or diminishes the sense of pain.

22. \_\_\_\_\_ a doctor who administers anesthesia.
23. \_\_\_\_\_ removal of extra skin at the head of a penis.
24. \_\_\_\_\_ tightening of uterine muscles that help move the baby from the uterus to and through the birth canal.
25. \_\_\_\_\_ refers to the size of the round opening of the cervix. An opening ten centimeters or large enough for five fingers to enter is full dilation.
26. \_\_\_\_\_ when the length of the cervical canal is shortened because it is thinning out to prepare for birth.
27. \_\_\_\_\_ cutting of perineum during birth to prevent skin from tearing and make the delivery easier.
28. \_\_\_\_\_ returning of uterus from pregnancy size to non-pregnant state.
29. \_\_\_\_\_ same as miscarriage or natural birth of fetus before full term.
30. \_\_\_\_\_ the physician trained to care for children up to the age of adolescence.
31. \_\_\_\_\_ attached at the end of the umbilical cord to the uterine wall to transport nourishment from mother to baby.
32. \_\_\_\_\_ anesthetic given through buttocks to numb the pelvic area (includes nerve fibers in the external organs, perineum, vagina, and vulva).
33. \_\_\_\_\_ relieves all sensation of pain.
34. \_\_\_\_\_ when warm water is inserted into the rectum to remove the solid waste.
35. \_\_\_\_\_ a heavy vaginal discharge after the birth of the baby. The average flow is 21 days.
36. \_\_\_\_\_ given in a space at the base of the back bone. It numbs the lower back and buttocks.

37. \_\_\_\_\_ physician who is trained in female concerns but also gives prenatal care, delivers babies, and gives postnatal care.
38. \_\_\_\_\_ movements of the baby into head down position at the end of pregnancy providing relief of pressure on the rib cage.
39. \_\_\_\_\_ when a baby is born bottom or feet first.
40. \_\_\_\_\_ the cervix dilates from 3-7 cm, contractions 2-3 minutes apart.
41. \_\_\_\_\_ skin between vaginal opening and the anus which is cut during delivery.
42. \_\_\_\_\_ when the baby's head can be seen at the vaginal opening and is ready to be born.
43. \_\_\_\_\_ a baby born before developing full term and weighing under 5 1/2 pounds.
44. \_\_\_\_\_ natural birth of fetus before it is developed fully, not full term.
45. \_\_\_\_\_ a visual reading of the newborn that checks color, respiration, reflex irritability, pulse, and muscle tone.
46. \_\_\_\_\_ the work of the mother and baby to help the baby be delivered.